



ETHICS and DOPING Focus on cobalt

Dr Brieuc DE MOFFARTS DVM, MSc, PhD



Dr Marie-Capucine DUPUIS TRICAUD DVM, MCMVS, PhD



Cobalt inquiry: Horses given 'unknown' injections

December 21, 2015



Horse racing trainers resigned to bans for cobalt

THE AUSTRALIAN | JANUARY 20, 2016 12:00AM



IS COBALT A KILLER IN HORSES?


Australian Vets Condemn Cobalt Misuse in Horses

Ethics is a question of recognized values, for example:

Health


Well-Being

Performance




Public point of view

Professional point of view




• Cobalt:

- Mineral element component of B12 vitamin (cobalamin)
- Essential for red blood cell formation
- Induces numerous genes expressions, but will rapidly be toxic
- Cobalamin is not able to enhance performance except if deficiency is diagnosed



- Nutritionally a horse needs
 - 1mg/day at rest
 - 7mg/day in full training
- Legally EuroLex indicates a maximal daily intake of 10mg in additional feeds.






 But cobalt in equine also supports performance and can be detrimental to health and well-being



40mg is a subtoxic dose



- Extended level of cobalt in equine is considered doping
- A level of 100ng/ml of cobalt in urine seems to be the upper limit in Europe.
- Additionally 25ηg/ml of cobalt in plasma is also a relevant threshold (US).





- Daily needs for a horse in training: 7 mg
- TWYDIL RACING: 3.5mg/day
- TWYDIL HEMATINIC: 5 to 6 mg/day
- TWYDIL HEMOPAR: 0.5mg/day
- 2 forms: Cobalamin (vit B12) or inorganic sources of Co++












Effet of TWYDIL HEMATINIC on blood parameters

- Evaluation of the TWYDIL HEMATINIC components bioavaibility
- 2. Evaluation on healthy horses of the impact of the supplement on red blood cells regenerative mechanism

Effet of TWYDIL HEMATINIC on blood parameters




1. Evaluation of the TWYDIL HEMATINIC components bioavaibility



Effet of TWYDIL HEMATINIC on blood parameters




1. Evaluation of the TWYDIL HEMATINIC components bioavaibility



Effet of TWYDIL HEMATINIC on blood parameters



1. Evaluation of the TWYDIL HEMATINIC components bioavaibility



10 days of TWYDIL HEMATINIC (Co++ 6.8mg) allow a significant increase of red blood cells constitutive elements and protective factors.







Effet of TWYDIL HEMATINIC on blood parameters



2. Impact on red blood cells regeneration










Effet of TWYDIL HEMATINIC on blood parameters




2. Impact on red blood cells regeneration

Haematocrit variation in supplemented group (%)



Folic acid concentration variation in supplemented group (ng.mL-1)

Haematocrit variation in placebo group (%)



Folic acid concentration variation in placebo group (ηg.mL-1)

- A 10 day supplementation with TWYDIL HEMATINIC seems to support red blood cells regenerative mechanism
- in healthy and aneamic horses without any risk of toxicity or even congestion
- this efficacy is mostly linked to the folic acid intake.





Effect of TWYDIL program / Cobalt

Purposes:

Effet of TWYDIL program on cobalt concentration

Evaluation of urine and plasmatic level of cobalt during 10 days of TWYDIL RACING and TWYDIL HEMATINIC administration

corresponding to a total amount of 11 mg of cobalt per day






Effet on concentration of cobalt





 Impact on urine concentration of 11mg of cobalt after 10 days of administration



Time (h) after last administration


Effet on concentration of cobalt





2. Impact on plasmatic concentration of 11mg of cobalt during 10 days of administration

(US Threshold: 25 ηg/ml)



Time (days) of administration

Conclusion:

Effet on concentration of cobalt

- No cumulative effect of cobalt orally absorbed in urine
- Possibility of cumulative effect in plasma
- 72h are sufficient in order to eliminate totally the cobalt in the urine

Estimation: 20 mg/day of orally absorbed cobalt is the limit beyond which there is a risk to exceed 25 ηg of cobalt per ml of plasma, for a probable duration over 72hours.

CONCLUSION

- Even with a legal daily dose of cobalt into the feed, a risk of positive case during the 10 hours following the administration is present.
- Respect dose and duration

